



Imaging of airway remodelling

Learn more about the benefits, limitations and clinical correlates of imaging techniques available for the assessment of airway remodelling in asthma



EpiCentral
UNDERSTANDING THE CENTRAL ROLE OF THE
EPITHELIUM IN SEVERE ASTHMA AND BEYOND

CT assessment of airway remodelling

Benefits¹⁻⁶

- Gold standard in pulmonary imaging
- Airway wall thickness measurements are consistent with histological examinations
- Simple to measure lung parenchymal density and gas trapping
- Allows identification and quantification of mucus plugging


Limitations^{1,2,6}

- Complex and potentially difficult to measure airway dimensions
- Low precision for measuring small airways of <1–2 mm diameter
- Cannot distinguish which specific components of the airway wall are thickened in airway remodelling

Clinical correlates^{3,7-12}

- CT-assessed remodelling has been found to correlate positively with asthma severity in adult patients
- CT assessment of gas trapping is associated with asthma severity and airway hyperresponsiveness
- Patients with increased WA% and gas trapping on CT scans may be more likely to exhibit neutrophilic inflammation
- Mucus plugging shown on CT is associated with sputum eosinophilia

Bronchial wall thickening in a patient with severe asthma as assessed by inspiratory CT¹³



CT, computed tomography; WA%, wall area percentage

1. King GG, et al. Eur Respir Rev 2019;28:180111; 2. Dournes G, Laurent F. Pulm Med 2012;2012:670414; 3. Dunican EM, et al. J Clin Invest 2018;128:997–1009; 4. Trivedi A, et al. J Allergy Clin Immunol 2017;139:1–10; 5. Stewart NJ, et al. Br J Radiol 2022;95:20210207; 6. de Jong PA, et al. Eur Respir J 2005;26:140–152; 7. Aysola RS, et al. Chest 2008;134:1183–1191; 8. Niimi A, et al. Am J Crit Care Med 2000;162:1518–1523; 9. Little SA, et al. Thorax 2002;57:247–253; 10. Busacker A, et al. Chest 2009;135:48–56; 11. Ueda T, et al. J Allergy Clin Immunol 2006;118:1019–1025; 12. Gupta S, et al. Thorax 2010;65:775–781; 13. van den Bosch WB, et al. Eur Respir Rev 2021;30:200186

Z4-55653; date of preparation: June 2023. © 2023 AstraZeneca. All Rights Reserved. This information is intended for healthcare professionals only. EpiCentral is sponsored and developed by AstraZeneca.

MRI assessment of airway remodelling

Benefits¹⁻⁵

- High spatial and temporal resolution of ventilation defects, which reflect airway narrowing
- Allows longitudinal monitoring of disease with avoidance of exposure of patient to ionising radiation
- Can provide information not captured by pulmonary function tests


Limitations^{1,4,6}

- Airway wall thickness cannot be measured
- Slightly reduced spatial resolution compared with CT scans

Clinical correlates^{1-3,7-10}

- MRI ventilation defects are associated with asthma severity and exacerbation risk
- Ventilation defects predict long-term FEV₁ reversibility in mild-to-moderate asthma
- Ventilation defects may also be associated with eosinophilia and poor control of eosinophilic inflammation
- Regions of air trapping and mucus plugging on CT overlap with MRI-assessed ventilation defects

Hyperpolarised noble gas MRI static ventilation images of patients with mild, moderate and severe asthma^{2,11*}



*As defined by GINA guidelines¹¹

CT, computed tomography; FEV₁, forced expiratory volume in 1 second; GINA, Global Initiative for Asthma; MRI, magnetic resonance imaging

1. King GG, et al. Eur Respir Rev 2019;28:180111; 2. Kooner HK, et al. Respirology 2022;27:114–133; 3. Stewart NJ, et al. Br J Radiol 2022;95:20210207; 4. Trivedi A, et al. J Allergy Clin Immunol 2017;139:1–10; 5. Petousi N, et al. Thorax 2019;74:797–805; 6. de Jong PA, et al. Eur Respir J 2005;26:140–152; 7. Mummy DG, et al. J Allergy Clin Immunol 2018;141:1140–1141; 8. Altes TA, et al. J Allergy Clin Immunol 2016;137:789–796; 9. Svenningsen S, et al. Am J Respir Crit Care Med 2018;197:876–884; 10. Eddy RL, et al. Radiology 2019;293:212–220; 11. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. 2023. Available from: <https://ginasthma.org/wp-content/uploads/2023/05/GINA-2023-Full-Report-2023-WMS.pdf> (Accessed 7 June 2023)

Z4-55653; date of preparation: June 2023. © 2023 AstraZeneca. All Rights Reserved. This information is intended for healthcare professionals only. EpiCentral is sponsored and developed by AstraZeneca.

Benefits¹⁻³

- More sensitive method to study bronchial wall thickness than HRCT
- Capability to discriminate between individual layers of the airways; allows 3–5 layers to be distinguished
- Access to airways as small as 4 mm


Limitations^{1,2}

- Requires bronchoscopy, which carries a risk of bronchospasm
- Standards have not yet been established

Clinical correlates^{4,5}

- Thickening of bronchial walls L₁, L₂ and L₃₋₅, as measured by EBUS, is associated with severe asthma
- PC₂₀, a measure of airway hyperresponsiveness, negatively correlates with the thickness of the second airway wall layer in patients with asthma

Measurement of bronchial wall thickness using HRCT and EBUS¹



EBUS, endobronchial ultrasound; FEV₁, forced expiratory volume in 1 second; HRCT, high-resolution computed tomography; L, layer; PC₂₀, provocation concentration of methacholine causing a 20% fall in FEV₁

1. Gorska K, et al. Respir Med 2016;117:131–138; 2. Trivedi A, et al. J Allergy Clin Immunol 2017;139:1–10; 3. Manso L, et al. Allergol Immunopathol (Madr) 2012;40:108–116;

4. Soja J, et al. Pol Arch Med Wewn 2015;125:659–665; 5. Kita T, et al. J Bronchology Interv Pulmonol 2010;17:301–306

Z4-55653; date of preparation: June 2023. © 2023 AstraZeneca. All Rights Reserved. This information is intended for healthcare professionals only. EpiCentral is sponsored and developed by AstraZeneca.